

About This Guide

This Guide provides a high-level overview of Patient Tasks in Care360. The overview is meant to provide guidance for you, your practice electronic health record (EHR) champion, or IT staff.

Experienced users are the main focus of this Guide; not every step is included in the instructions, and this is not a replacement for training from Care360. There are several ways to approach each workflow in Care360. This Guide highlights one workflow; however, you may be familiar with an alternative approach. Please note that this Guide was created based upon Care360 version 2016.2. Screens and features may change as new software versions are released.

INDICATIONS AND USAGE

- PRALUENT is a PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9) inhibitor antibody indicated as adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or clinical atherosclerotic cardiovascular disease, who require additional lowering of LDL-C
- The effect of PRALUENT on cardiovascular morbidity and mortality has not been determined

IMPORTANT SAFETY INFORMATION

- PRALUENT is contraindicated in patients with a history of a serious hypersensitivity reaction to PRALUENT. Reactions have included hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization

Please see accompanying full [Prescribing Information](#)



Using Patient Tasks

Clinical decision support (CDS) tools such as Patient Tasks provide clinicians and staff with patient-specific information, intelligently filtered and presented at appropriate times to help improve the delivery of care. Patient Tasks can assist with identifying gaps-in-care and may enhance patient outcomes and improve care consistency—a benefit to both providers and patients.

Patient Tasks can be created to alert health care professionals (HCPs) to consider PRALUENT® therapy for appropriate patients during the visit.

Factors That May Impact Patient Tasks

The display of Patient Tasks may be impacted by the clinical data available in the EHR; for example, if lab results are saved in the EHR as a PDF file and not available for use as 'data' to be queried. Additionally in those cases where lab results are received from multiple laboratories, it may be necessary to select each lab's order codes to enable all appropriate patients to appear on the patient list. Also, medications prescribed before the EHR was implemented might not be included in a patient's medications list. These information gaps can limit the number of patients where Patient Tasks are displayed.

Patient Tasks can help identify whether the patient is a candidate for PRALUENT treatment based on clinical appropriateness and payer utilization management criteria via:

- Reminders during the patient exam workflow, such as ordering lab tests or other diagnostics, adjusting or adding medications to a patient's treatment plan, or providing patient education on LDL-C goals
- Identification of patients who, for example, have clinical atherosclerotic cardiovascular disease (ASCVD) and are on maximally tolerated statin therapy atorvastatin 40 mg/day and having elevated LDL-C ≥ 70 mg/dL or ≥ 100 mg/dL, depending on insurance

IMPORTANT SAFETY INFORMATION

- Hypersensitivity reactions (e.g., pruritus, rash, urticaria), including some serious events (e.g., hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization), have been reported with PRALUENT treatment. If signs or symptoms of serious allergic reactions occur, discontinue treatment with PRALUENT, treat according to the standard of care, and monitor until signs and symptoms resolve

Please see accompanying full [Prescribing Information](#)



Praluent[®]
(alirocumab) Injection 75mg/mL
150mg/mL
Redefining Possible

Reminders: Using Patient Tasks

Care360 enables the setup of reminders (Patient Tasks) which are displayed when the patient's chart is accessed. Patient Tasks are based on certain criteria. Available criteria include diagnosis, current and prior medications, lab values, and other clinical or patient demographic information.

For example, Patient Tasks may be created to alert the HCP of patients with uncontrolled LDL-C who are clinically appropriate and approvable for PRALUENT therapy based upon the approved Indication.

Patient Tasks can be created manually based on a patient summary report, which can be created listing all patients meeting specific criteria.

The following steps illustrate how to create patient tasks to identify patients who may be candidates for treatment with PRALUENT.

To Create a Patient Task from the Search Results List:

- Using the EHR patient identification tip sheet, create a Patient Summary Report (eg, diagnosis ASCVD, atorvastatin 40 mg/day, LDL-C ≥ 70 mg/dL or ≥ 100 mg/dL, depending on insurance)

| Test Name or Pharmacy Name of Allergy or Diagnosis Description | Results or Medications | Values or Units or Codes or Reactions | Patient Name | Patient ID | Birth Date | Lab ID | Lab Collected or Med Issued on or Allergy Modified or Diagnosis Date | Provider | Status | Out of Range |
|--|------------------------|---------------------------------------|--------------|------------|------------|--------|--|---------------|--------|--------------|
| | atorvastatin | 40 mg | | 114 | 10/31/1951 | | 10/27/2016 | William Folds | Active | |
| | pravastatin | 80 mg | | 328 | 07/03/1928 | | 10/26/2016 | William Folds | Active | |
| | simvastatin | 40 mg | | 2279 | 08/23/1956 | | 10/21/2016 | Thomas Jolly | Active | |
| | rosuvastatin | 20 mg | | 111 | 10/15/1935 | | 10/19/2016 | William Folds | Active | |
| | atorvastatin | 40 mg | | 854 | 07/15/1940 | | 10/18/2016 | Thomas Jolly | Active | |
| | simvastatin | 40 mg | | 854 | 07/15/1940 | | 10/17/2016 | Thomas Jolly | Active | |

IMPORTANT SAFETY INFORMATION


- The most commonly occurring adverse reactions ($\geq 5\%$ of patients treated with PRALUENT and occurring more frequently than with placebo) are nasopharyngitis, injection site reactions, and influenza

Please see accompanying full [Prescribing Information](#)



Reminders: Using Patient Tasks

2. Click on the **patient name** in Informatics Search Results to navigate to the Patient Summary screen
3. From the Patient Summary, Patient Tasks panel, click **Add**



4. Enter appropriate Task details
 - Select **General Task** from the Task Type dropdown
 - Select **appropriate HCP** in the **Assigned To field**
 - Select the **Due Date** to correspond to the patient's next appointment
 - Add appropriate message, eg, "**Consider patient for treatment with PRALUENT**"
5. Click **Save**

IMPORTANT SAFETY INFORMATION

- Local injection site reactions including erythema/redness, itching, swelling, and pain/tenderness were reported more frequently in patients treated with PRALUENT 75 mg and/or 150 mg every 2 weeks (7.2% versus 5.1% for PRALUENT and placebo, respectively). Few patients discontinued treatment because of these reactions (0.2% versus 0.4% for PRALUENT and placebo, respectively), but patients receiving PRALUENT had a greater number of injection site reactions, had more reports of associated symptoms, and had reactions of longer average duration than patients receiving placebo

Please see accompanying full [Prescribing Information](#)



To View Patient Tasks:

Tasks can be viewed from the HCP's Message Center.

The screenshot shows the Care360 Message Center interface. At the top, there are navigation tabs for 'Message Center', 'Patient', 'Lab Orders', and 'Reports'. Below this, there are sections for 'Find Lab Results', 'New Lab Results' (6), 'New Clinical Documents' (0), 'Failed Fax Messages' (41), and 'Results Pending Release' (12). A 'Messages' section indicates there are no new messages. A 'Tasks' table is visible at the bottom of the screenshot.

| Type | Subject | Due Date | Patient Name | Assigned User | Reserved Date |
|---------|--|------------|--------------|---------------|---------------------|
| General | Consider patient for treatment intensification with PRALUENT | 11/01/2016 | test, test | | 11/01/2016 12:28 PM |

Tasks are also displayed from the Patient Summary screen when accessing a patient chart.

The screenshot shows the Patient Summary screen. The 'Patient Tasks' section has an 'Add' link and a 'View All' link with an 'X' icon. Below it, a task is listed: '11/01/2016 Consider patient for treatment with PRALUENT' with a 'General' category. The 'Medications (Active)' section has links for 'Write a Prescription' and 'Reconcile', and a 'View All' link with an 'X' icon. Below it, a medication is listed: 'rosuvastatin' with a dosage of '20 mg tablet; TAKE 1 TABLET DAILY; Dispense 30...'.

IMPORTANT SAFETY INFORMATION

- The once-monthly (Q4W) 300 mg dosing regimen had a higher rate of local injection site reactions as compared to PRALUENT 75 mg Q2W or placebo (16.6%, 9.6%, and 7.9%, respectively) in a trial in which all patients received an injection of drug or placebo every 2 weeks to maintain the blind. The discontinuation rate due to injection site reactions was 0.7% in the 300 mg Q4W arm and 0% in the other 2 arms

Please see accompanying full [Prescribing Information](#)



INDICATIONS AND USAGE

- PRALUENT® (alirocumab) is a PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9) inhibitor antibody indicated as adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or clinical atherosclerotic cardiovascular disease, who require additional lowering of LDL-C
- The effect of PRALUENT on cardiovascular morbidity and mortality has not been determined

IMPORTANT SAFETY INFORMATION

- PRALUENT is contraindicated in patients with a history of a serious hypersensitivity reaction to PRALUENT. Reactions have included hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization
- Hypersensitivity reactions (e.g., pruritus, rash, urticaria), including some serious events (e.g., hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization), have been reported with PRALUENT treatment. If signs or symptoms of serious allergic reactions occur, discontinue treatment with PRALUENT, treat according to the standard of care, and monitor until signs and symptoms resolve
- The most commonly occurring adverse reactions ($\geq 5\%$ of patients treated with PRALUENT and occurring more frequently than with placebo) are nasopharyngitis, injection site reactions, and influenza
- Local injection site reactions including erythema/redness, itching, swelling, and pain/tenderness were reported more frequently in patients treated with PRALUENT 75 mg and/or 150 mg every 2 weeks (7.2% versus 5.1% for PRALUENT and placebo, respectively). Few patients discontinued treatment because of these reactions (0.2% versus 0.4% for PRALUENT and placebo, respectively), but patients receiving PRALUENT had a greater number of injection site reactions, had more reports of associated symptoms, and had reactions of longer average duration than patients receiving placebo
- The once-monthly (Q4W) 300 mg dosing regimen had a higher rate of local injection site reactions as compared to PRALUENT 75 mg Q2W or placebo (16.6%, 9.6%, and 7.9%, respectively) in a trial in which all patients received an injection of drug or placebo every 2 weeks to maintain the blind. The discontinuation rate due to injection site reactions was 0.7% in the 300 mg Q4W arm and 0% in the other 2 arms
- Neurocognitive events were reported in 0.8% of patients treated with PRALUENT and 0.7% of patients treated with placebo. Confusion or memory impairment were reported more frequently by those treated with PRALUENT (0.2% for each) than in those treated with placebo ($<0.1\%$ for each)
- Liver-related disorders (primarily related to abnormalities in liver enzymes) were reported in 2.5% of patients treated with PRALUENT and 1.8% of patients treated with placebo, leading to treatment discontinuation in 0.4% and 0.2% of patients, respectively. Increases in serum transaminases to greater than 3 times the upper limit of normal occurred in 1.7% of patients treated with PRALUENT and 1.4% of patients treated with placebo
- The most common adverse reactions leading to treatment discontinuation in patients treated with PRALUENT were allergic reactions (0.6% versus 0.2% for PRALUENT and placebo, respectively) and elevated liver enzymes (0.3% versus $<0.1\%$)
- PRALUENT is a human monoclonal antibody. As with all therapeutic proteins, there is a potential for immunogenicity with PRALUENT

Please see accompanying full [Prescribing Information](#)



©2017 Sanofi and Regeneron Pharmaceuticals, Inc.
All rights reserved. November 2017 SAUS.PRL.17.11.9049

