

About This Guide

This Guide provides a high-level overview of Reminder Tasks in Practice Fusion. The overview is meant to provide guidance for you, your practice electronic health record (EHR) champion, or IT staff.

Experienced users are the main focus of this Guide; not every step is included in the instructions, and this is not a replacement for training from Practice Fusion. There are several ways to approach each workflow in Practice Fusion. This Guide highlights one workflow; however, you may be familiar with an alternative approach. Please note that this Guide was created based upon Practice Fusion version 3.6.5.56.09.910. Screens and features may change as new software versions are released.

INDICATIONS AND USAGE

- PRALUENT is a PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9) inhibitor antibody indicated as adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or clinical atherosclerotic cardiovascular disease, who require additional lowering of LDL-C
- The effect of PRALUENT on cardiovascular morbidity and mortality has not been determined

IMPORTANT SAFETY INFORMATION

- PRALUENT is contraindicated in patients with a history of a serious hypersensitivity reaction to PRALUENT. Reactions have included hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization

Please see accompanying full [Prescribing Information](#)



Using Reminder Tasks

Clinical decision support (CDS) tools such as Reminder Tasks provide clinicians and staff with patient-specific information, intelligently filtered and presented at appropriate times to help improve the delivery of care. Reminder Tasks can assist with identifying gaps-in-care and may enhance patient outcomes and improve care consistency—a benefit to both providers and patients.

Reminder Tasks can be created to alert health care professionals (HCPs) to consider PRALUENT therapy for appropriate patients during the visit.

Factors That May Impact Reminder Tasks

The display of Reminder Tasks may be impacted by the clinical data available in the EHR; for example, if lab results are saved in the EHR as a PDF file and not available for use as “data” to be queried. Additionally, in those cases where lab results are received from multiple laboratories, it may be necessary to select each lab’s order codes to enable all appropriate patients to display Reminder Tasks. Also, medications prescribed before the EHR was implemented might not be included in a patient’s medications list. These information gaps can limit the number of patients where alerts are displayed.

Reminder Tasks can help identify whether the patient is a candidate for PRALUENT treatment based on clinical appropriateness and payer utilization management criteria via:

- Reminders during the patient exam workflow, such as ordering lab tests or other diagnostics, adjusting or adding medications to a patient’s treatment plan, or providing patient education on LDL-C goals
- Identification of patients who, for example, have clinical atherosclerotic cardiovascular disease (ASCVD) and are on maximally tolerated statin therapy atorvastatin 40 mg/day and having elevated LDL-C ≥ 70 mg/dL or ≥ 100 mg/dL, depending on insurance

IMPORTANT SAFETY INFORMATION

- Hypersensitivity reactions (e.g., pruritus, rash, urticaria), including some serious events (e.g., hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization), have been reported with PRALUENT treatment. If signs or symptoms of serious allergic reactions occur, discontinue treatment with PRALUENT, treat according to the standard of care, and monitor until signs and symptoms resolve

Please see accompanying full [Prescribing Information](#)



Reminders: Using Reports and Tasks

Practice Fusion enables reminders (Tasks) which are displayed when the patient's chart is accessed.

For example, a task may be created to alert the HCP of patients with uncontrolled LDL-C who are clinically appropriate and approvable for PRALUENT therapy based upon the approved Indication.

Tasks can be created manually based on a patient list report that can be created listing all patients meeting specific criteria. Criteria for PRALUENT would include clinical ASCVD, being on maximally tolerated statin therapy (eg, atorvastatin 40 mg/day) and having elevated LDL-C (eg, ≥ 70 mg/dL or ≥ 100 mg/dL), depending on insurance.

The following steps illustrate how to create Tasks to identify patients who may be candidates for treatment with PRALUENT.

To Add a Reminder Task:

Use the EHR tip sheet to identify patients that may be appropriate for PRALUENT. For each patient on the resulting list, complete these steps:

1. From the Patient List on the screen, click on the **patient name** to access the patient chart
2. Click on **Actions, Create Task**
3. Enter Task **Details** such as "Consider patient for treatment with PRALUENT"
4. Add a **Reminder Date** to correspond with the patient's next appointment
5. Assign the Task to the HCP
6. Click **Save**

IMPORTANT SAFETY INFORMATION

- The most commonly occurring adverse reactions ($\geq 5\%$ of patients treated with PRALUENT and occurring more frequently than with placebo) are nasopharyngitis, injection site reactions, and influenza

Please see accompanying full [Prescribing Information](#)



To View a Reminder

Manually created Tasks display as Reminders, and can be viewed from the Task section, Reminders tab. Reminders are displayed in date order.

The screenshot shows the 'Tasks' section in Practice Fusion. At the top, there are tabs for 'All tasks (2)', 'My task (4)', 'Unassigned tasks (6)', 'Lab results (3)', and 'Reminders (7)'. Below the tabs are filters for 'Reminder tasks', 'Incomplete', 'Current', and 'All providers'. There are also buttons for 'Connect Labs', 'Connect imaging centers', and 'New task'. The main table has columns for 'DATE', 'TASK', 'PATIENT', 'DETAILS', and 'ACTIONS'. A single reminder is listed for 'Yesterday' on '12/20/2016' for a 'Reminder' task for 'Zelda Testpatient' (62 yrs. F. DOB 08/12/1954). The details are 'Consider patient for treatment with PRALUENT'. An 'Actions' dropdown menu is visible next to the task.

DATE	TASK	PATIENT	DETAILS	ACTIONS
Yesterday				
12/20/2016	Reminder	Zelda Testpatient 62 yrs. F. DOB 08/12/1954	Consider patient for treatment with PRALUENT	View

IMPORTANT SAFETY INFORMATION

- Local injection site reactions including erythema/redness, itching, swelling, and pain/tenderness were reported more frequently in patients treated with PRALUENT 75 mg and/or 150 mg every 2 weeks (7.2% versus 5.1% for PRALUENT and placebo, respectively). Few patients discontinued treatment because of these reactions (0.2% versus 0.4% for PRALUENT and placebo, respectively), but patients receiving PRALUENT had a greater number of injection site reactions, had more reports of associated symptoms, and had reactions of longer average duration than patients receiving placebo

Please see accompanying full [Prescribing Information](#)



INDICATIONS AND USAGE

- PRALUENT® (alirocumab) is a PCSK9 (Proprotein Convertase Subtilisin/Kexin Type 9) inhibitor antibody indicated as adjunct to diet and maximally tolerated statin therapy for the treatment of adults with heterozygous familial hypercholesterolemia or clinical atherosclerotic cardiovascular disease, who require additional lowering of LDL-C
- The effect of PRALUENT on cardiovascular morbidity and mortality has not been determined

IMPORTANT SAFETY INFORMATION

- PRALUENT is contraindicated in patients with a history of a serious hypersensitivity reaction to PRALUENT. Reactions have included hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization
- Hypersensitivity reactions (e.g., pruritus, rash, urticaria), including some serious events (e.g., hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization), have been reported with PRALUENT treatment. If signs or symptoms of serious allergic reactions occur, discontinue treatment with PRALUENT, treat according to the standard of care, and monitor until signs and symptoms resolve
- The most commonly occurring adverse reactions ($\geq 5\%$ of patients treated with PRALUENT and occurring more frequently than with placebo) are nasopharyngitis, injection site reactions, and influenza
- Local injection site reactions including erythema/redness, itching, swelling, and pain/tenderness were reported more frequently in patients treated with PRALUENT 75 mg and/or 150 mg every 2 weeks (7.2% versus 5.1% for PRALUENT and placebo, respectively). Few patients discontinued treatment because of these reactions (0.2% versus 0.4% for PRALUENT and placebo, respectively), but patients receiving PRALUENT had a greater number of injection site reactions, had more reports of associated symptoms, and had reactions of longer average duration than patients receiving placebo
- The once-monthly (Q4W) 300 mg dosing regimen had a higher rate of local injection site reactions as compared to PRALUENT 75 mg Q2W or placebo (16.6%, 9.6%, and 7.9%, respectively) in a trial in which all patients received an injection of drug or placebo every 2 weeks to maintain the blind. The discontinuation rate due to injection site reactions was 0.7% in the 300 mg Q4W arm and 0% in the other 2 arms
- Neurocognitive events were reported in 0.8% of patients treated with PRALUENT and 0.7% of patients treated with placebo. Confusion or memory impairment were reported more frequently by those treated with PRALUENT (0.2% for each) than in those treated with placebo ($<0.1\%$ for each)
- Liver-related disorders (primarily related to abnormalities in liver enzymes) were reported in 2.5% of patients treated with PRALUENT and 1.8% of patients treated with placebo, leading to treatment discontinuation in 0.4% and 0.2% of patients, respectively. Increases in serum transaminases to greater than 3 times the upper limit of normal occurred in 1.7% of patients treated with PRALUENT and 1.4% of patients treated with placebo
- The most common adverse reactions leading to treatment discontinuation in patients treated with PRALUENT were allergic reactions (0.6% versus 0.2% for PRALUENT and placebo, respectively) and elevated liver enzymes (0.3% versus $<0.1\%$)
- PRALUENT is a human monoclonal antibody. As with all therapeutic proteins, there is a potential for immunogenicity with PRALUENT

Please see accompanying full [Prescribing Information](#)



©2017 Sanofi and Regeneron Pharmaceuticals, Inc.
All rights reserved. November 2017 SAUS.PRL.17.11.9040

