

About This Guide

This Guide provides a high-level overview of Clinical Decision Support (CDS) Rules in Amazing Charts. The overview is meant to provide guidance for you, your practice electronic health record (EHR) champion, or IT staff.

Experienced users are the main focus of this Guide; not every step is included in the instructions, and this is not a replacement for training from Amazing Charts. There are several ways to approach each workflow in Amazing Charts. This Guide highlights one workflow; however, you may be familiar with an alternative approach. Please note that this Guide was created based upon Amazing Charts version 9.3. Screens and features may change as new software versions are released.

INDICATIONS AND USAGE

PRALUENT (alirocumab) is indicated:

- to reduce the risk of myocardial infarction, stroke, and unstable angina requiring hospitalization in adults with established cardiovascular disease
- as an adjunct to diet, alone or in combination with other lipid-lowering therapies (e.g., statins, ezetimibe), for the treatment of adults with primary hyperlipidemia (including heterozygous familial hypercholesterolemia) to reduce low-density lipoprotein cholesterol (LDL-C)

IMPORTANT SAFETY INFORMATION

- PRALUENT is contraindicated in patients with a history of a serious hypersensitivity reaction to PRALUENT, including hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization

Please see accompanying full [Prescribing Information](#)



Using CDS Rules

Clinical decision support (CDS) tools such as CDS Rules provide clinicians and staff with patient-specific information, intelligently filtered and presented at appropriate times to help improve the delivery of care. CDS Rules can assist with identifying gaps-in-care and may enhance patient outcomes and improve care consistency—a benefit to both providers and patients.

Factors That May Impact CDS Rules

The display of CDS Rules may be impacted by the clinical data available in the EHR; for example, if lab results are saved in the EHR as a PDF file and not available for use as “data” to be queried. Additionally, in those cases where lab results are received from multiple laboratories, it may be necessary to select each lab’s order codes to enable all appropriate patients to display reminders.

The query criteria should consider active patients only (not deceased or inactive, as determined by the practice). Also, medications prescribed before the EHR was implemented might not be included in a patient’s medications list. These information gaps can limit the number of patients where CDS Rules are displayed.

CDS Rules can be created to alert health care professionals (HCPs) to consider PRALUENT therapy for appropriate patients during the visit. CDS Rules can help identify whether the patient is a candidate for PRALUENT treatment based on clinical appropriateness and payer utilization management criteria via:

- CDS Rules during the patient exam workflow, such as ordering lab tests or other diagnostics, adjusting or adding medications to a patient’s treatment plan, or providing patient education on LDL-C goals
- Identification of patients who, for example, have established cardiovascular disease (eg, myocardial infarction, stroke or unstable angina requiring hospitalization) and may also be on maximally tolerated statin therapy (eg, atorvastatin 40 mg/day), and having elevated LDL-C ≥ 70 mg/dL or ≥ 100 mg/dL, depending on insurance

IMPORTANT SAFETY INFORMATION

- Hypersensitivity reactions (e.g., pruritus, rash, urticaria), including some serious events (e.g., hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization), have been reported with PRALUENT treatment. If signs or symptoms of serious allergic reactions occur, discontinue treatment with PRALUENT, treat according to the standard of care, and monitor until signs and symptoms resolve

Please see accompanying full [Prescribing Information](#)



Praluent[®]
(alirocumab) Injection 75mg/mL
150mg/mL
Redefining Possible

Using Clinical Decision Support Rules

Amazing Charts enables the setup of reminders based on criteria set in Clinical Decision Support Rules. Available criteria include diagnosis, current and prior medications, lab values, and other clinical or patient demographic information. The Clinical Decision Support reminder is displayed when the data in the patient chart meet the criteria.

For example, a reminder may be created to alert the HCP of patients with uncontrolled LDL-C who are clinically appropriate and approvable for PRALUENT therapy based upon the approved Indication.

The following steps illustrate how to create clinical decision support reminders to identify patients who may be candidates for treatment with PRALUENT.

To Create and Activate a Clinical Decision Support Rule:

Use the EHR tip sheet to help identify patients that may be appropriate for PRALUENT. For each patient on the resulting list, complete these steps:

1. Navigate to **Manage Decision Support Rules**
2. From the **Rule Details** tab, enter the Name and Type of rule to be created, eg,
 - Name: Candidate for Treatment with PRALUENT
 - Type: Recommendation
3. Add appropriate **Rule Details**, and click **Add**

Category	Name	Operator	Value
Diagnosis/RF (has at least one)	ICD-10 Code	=	E78.4
	ICD-10 Code	=	E78.0
	ICD-10 Code	=	E78.2
	ICD-10 Code	=	E78.5
	ICD-10 Code	=	E78.01
	ICD-10 Code	=	E78.00
Drug Classes (has at least one)	Drug Class	=	HMG-CoA reductase inhibitors
	Drug Class	=	Antilipemic Agent, HMG-CoA Reductase
	Lab Name	Starts With	LDL
Lab's (has at least one)	Lab Value	>	100

IMPORTANT SAFETY INFORMATION

- The most commonly occurring adverse reactions in clinical trials in primary hyperlipidemia (including heterozygous familial hypercholesterolemia (HeFH)) ($\geq 5\%$ of patients treated with PRALUENT and occurring more frequently than with placebo) are nasopharyngitis, injection site reactions, and influenza

Please see accompanying full [Prescribing Information](#)



Reminders: Using Clinical Decision Support Rules:

4. Select the **Conditions** tab to select criteria for the rule

Click **Edit Conditions** to edit Categories

Name: select Medications from the dropdown list (eg, atorvastatin, simvastatin)

Operator: select appropriate Operator

Value: enter value of each criterion; for example, diagnosis, drug class (statins), laboratory results (eg, LDL-C \geq 70 mg/dL or \geq 100 mg/dL)

Click **Add**

Click **Save** to activate the rule

IMPORTANT SAFETY INFORMATION

- The most commonly occurring adverse reactions in the cardiovascular outcomes trial (>5% of patients treated with PRALUENT and occurring more frequently than placebo) were non-cardiac chest pain, nasopharyngitis, and myalgia

Please see accompanying full [Prescribing Information](#)



Praluent[®]
(alirocumab) Injection 75mg/mL
150mg/mL
Redefining Possible

To View a Reminder Triggered by a Clinical Decision Support Rule:

Amazing Charts reminders are viewed from within the Most Recent Encounter (MRE).

Plan | CDS | Instructions Declines Patient Summary Patient Ed Given

Name	Last	Next
<Reminder message here>	—	—
	—	Today
	—	

Right click on a recommendation for more info, or to edit it

The Reminder can also be viewed from the CDS tab.

Decisions Support Due | Immunizations & Shots | Screenings & Tests | Injections non-Decision Support

Below is a list of this patient's Decisions Support recommendations

Show immunizations only.
 Also show items not yet due.
 Hide childhood immunizations

Name	Last	Next
<Reminder message here>		

Refresh

IMPORTANT SAFETY INFORMATION

- In the primary hyperlipidemia (including HeFH) clinical trials, local injection site reactions including erythema/redness, itching, swelling, and pain/tenderness were reported more frequently in patients treated with PRALUENT 75 mg and/or 150 mg every 2 weeks (7.2% versus 5.1% for PRALUENT and placebo, respectively). Few patients discontinued treatment because of these reactions (0.2% versus 0.4% for PRALUENT and placebo, respectively), but patients receiving PRALUENT had a greater number of injection site reactions, had more reports of associated symptoms, and had reactions of longer average duration than patients receiving placebo

Please see accompanying full [Prescribing Information](#)



INDICATIONS AND USAGE

PRALUENT (alirocumab) is indicated:

- to reduce the risk of myocardial infarction, stroke, and unstable angina requiring hospitalization in adults with established cardiovascular disease
- as an adjunct to diet, alone or in combination with other lipid-lowering therapies (e.g., statins, ezetimibe), for the treatment of adults with primary hyperlipidemia (including heterozygous familial hypercholesterolemia) to reduce low-density lipoprotein cholesterol (LDL-C)

IMPORTANT SAFETY INFORMATION

- PRALUENT is contraindicated in patients with a history of a serious hypersensitivity reaction to PRALUENT, including hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization
- Hypersensitivity reactions (e.g., pruritus, rash, urticaria), including some serious events (e.g., hypersensitivity vasculitis and hypersensitivity reactions requiring hospitalization), have been reported with PRALUENT treatment. If signs or symptoms of serious allergic reactions occur, discontinue treatment with PRALUENT, treat according to the standard of care, and monitor until signs and symptoms resolve
- The most commonly occurring adverse reactions in clinical trials in primary hyperlipidemia (including heterozygous familial hypercholesterolemia (HeFH)) ($\geq 5\%$ of patients treated with PRALUENT and occurring more frequently than with placebo) are nasopharyngitis, injection site reactions, and influenza
- The most commonly occurring adverse reactions in the cardiovascular outcomes trial ($>5\%$ of patients treated with PRALUENT and occurring more frequently than placebo) were non-cardiac chest pain, nasopharyngitis, and myalgia
- In the primary hyperlipidemia (including HeFH) clinical trials, local injection site reactions including erythema/redness, itching, swelling, and pain/tenderness were reported more frequently in patients treated with PRALUENT 75 mg and/or 150 mg every 2 weeks (7.2% versus 5.1% for PRALUENT and placebo, respectively). Few patients discontinued treatment because of these reactions (0.2% versus 0.4% for PRALUENT and placebo, respectively), but patients receiving PRALUENT had a greater number of injection site reactions, had more reports of associated symptoms, and had reactions of longer average duration than patients receiving placebo
- The once-monthly (Q4W) 300mg dosing regimen had a higher rate of local injection site reactions as compared to PRALUENT 75mg Q2W or placebo (16.6%, 9.6%, and 7.9%, respectively) in a trial in which all patients received an injection of drug or placebo every 2 weeks to maintain the blind. The discontinuation rate due to injection site reactions was 0.7% in the 300 mg Q4W arm and 0% in the other 2 arms

Please see accompanying full [Prescribing Information](#)



Praluent[®]
(alirocumab) Injection 75mg/mL
150mg/mL
Redefining Possible

IMPORTANT SAFETY INFORMATION (*cont.*)

- In a cardiovascular outcomes trial, local injection site reactions were reported in 3.8% of patients treated with PRALUENT versus 2.1% patients treated with placebo, and led to permanent discontinuation in 0.3% of patients versus <0.1% of patients, respectively
- In the primary hyperlipidemia trials, liver-related disorders (primarily related to abnormalities in liver enzymes) were reported in 2.5% of patients treated with PRALUENT and 1.8% of patients treated with placebo, leading to treatment discontinuation in 0.4% and 0.2% of patients, respectively. Increases in serum transaminases to greater than 3 times the upper limit of normal occurred in 1.7% of patients treated with PRALUENT and 1.4% of patients treated with placebo
- In the primary hyperlipidemia trials, the most common adverse reactions leading to treatment discontinuation in patients treated with PRALUENT were allergic reactions (0.6% versus 0.2% for PRALUENT and placebo, respectively) and elevated liver enzymes (0.3% versus <0.1%)
- PRALUENT is a human monoclonal antibody. As with all therapeutic proteins, there is a potential for immunogenicity with PRALUENT

Please see accompanying full [Prescribing Information](#)

SANOFI  **REGENERON**

©2019 Sanofi and Regeneron Pharmaceuticals, Inc.
All rights reserved. 07/2019 SAUS.PRL.17.11.9053(1)


Praluent[®]
(alirocumab) Injection 75mg/mL
150mg/mL
Redefining Possible